arXiv:1202.2907v1 [cs.CR] 14 Feb 2012

The weight Enumerator of some irreducible cyclic codes

Yun Song , Zhihui Li *

College of Mathematics and Information Science, Shaanxi Normal University, Xi'an, 710062, P. R. China

---

Abstract

Irreducible cyclic codes are one of the largest known classes of block codes which have been investigated for a long time. However, their weight distributions are known only for a few cases. In this paper, a class of irreducible cyclic codes are studied and their weight distributions are determined. Moreover, all codewords of some irreducible cyclic codes are obtained through programming in order to explain their distributions. The number of distinct nonzero weights in these codes dealt with in this paper varies among 1,2,3,6,8.

Keywords: Gaussian periods; Period polynomial; Irreducible cyclic codes; Linear codes; Weight distributions; Weight enumerator

---

1. Introduction

Let $q = p^s$ and $r = q^m$, where s and m are positive integers, and p a prime. A linear $[n, k, d]$ code C over F_q is a k-dimensional subspace of F_q^n with minimum (Hamming) distance d. Let A_i denote the number of codewords in a code C with Hamming weight i. The weight enumerator of C is defined by

$$1 + A_1x + A_2x^2 + \cdots + A_nx^n. \qquad (1)$$

The sequence$(1, A_1, \ldots, A_n)$ is called the weight distribution of the code.

Let $N > 1$ be an integer dividing $r-1$, and put $n = \frac{(r-1)}{N}$. Let α be a primitive element of F_{q^m} and $\theta = \alpha^N$. The set

$$C(q,m,N) = \{\mathrm{Tr}_{r/q}(\beta), \mathrm{Tr}_{r/q}(\beta\theta), \cdots, \mathrm{Tr}_{r/q}(\beta\theta^{n-1}) | \beta \in F_r\} \qquad (2)$$

is called an irreducible cyclic code (ICC) over F_q, where Tr is the trace function from F_r onto F_q.

The determination of the weight enumerator of ICC is a fascinating problem, which provides some valuable informations about the performance for the code. Important contributions in this direction can be found in [1-3]. McEliece showed that the weights of an irreducible cyclic codes can be expressed as a linear combination of Gauss sums via the Fourier transform[4]. Schmidt and White gave a characterization of ICC with at most two weights[5]. Langevin investigated a class of ICC with at most three weights[6]. However, in general, determining the weight distribution of ICC is quite complicated. The following special cases were studied in the literature.

---

*Corresponding author:Zhihui Li.(snnulzh@yahoo.com.cn)

• When $N|(q^j+1)$ for some j being a divisor of $m/2$, then codes have two weights. These codes were discussed in [7].

• When $N=2$, the weight distribution of $C(r,n)$ was founded by Baumert and McEliece[7].

• When N is a prime with $N \equiv 3 (\mathrm{mod}\ 4)$ and $ord_q(N)=(N-1)/2$, the weight distribution was studied by Baumert and Mykkeltveit[8].

• When $2 \leq N \leq 4$, the number of distinct nonzero weights in the ICC dealt with in [9] varies between one and four.

The objective of this paper is to determine the weight enumerator of a class of ICC for all N with $5 \leq N \leq 8$. The number of distinct nonzero weights in these codes dealt with in this paper varies among 1,2,3,6,8.

2. Cyclotomic classes and period polynomials

To study the weight enumerator of the irreducible cyclic codes, we need to introduce cyclotomic classes and period polynomials[10].

Definition 2.1. *Let $r-1=nN$ for two positive integers $n>1$ and $N>1$, and let α be a fixed primitive element of F_r. Define $C_i^{(N,r)}=\alpha^i\langle\alpha^N\rangle$ for $i=0,1,\ldots,N-1$, where $\langle\alpha^N\rangle$ denotes the subgroup of F_r^*generated by $\langle\alpha^N\rangle$. The cosets $C_i^{(N,r)}$ are called the cyclotomic classes of order in F_r .*

Definition 2.2. *The Gaussian periods are defined by*

$$\eta_i^{(N,r)}=\sum_{x\in C_i^{(N,r)}}\chi(x), \qquad i=0,1,\ldots,N-1,$$

where χ is the canonical additive character of F_r.

To determine the weight distribution of the ICC, we need to compute the Gaussian periods.Unfortunally, Definition 2.2 is not enough to determine the values of the Gaussian periods η_i. To this end, we need to introduce the period polynomials.

Definition 2.3. *The period polynomials are defined by*

$$\psi_{(N,r)}(X)=\prod_{i=0}^{N-1}(X-\eta_i^{(N,r)}). \qquad (3)$$

It is known that $\psi_{(N,r)}(X)$ is a polynomial with integer coefficients [10]. *We will need the following four lemmas whose proofs can be found in* [10 − 11].

Lemma 2.4. *Let $N = 5$. We have the following results on the factorization of $\psi_{(5,r)}(X)$.*

(i) If $p \equiv 4(\mathrm{mod} 5)$, then

$$\psi_{(5,r)}(X) = \{ \begin{array}{ll} 5^{-5}(5X+1+4\sqrt{r})(5X+1-\sqrt{r}), & if sm/2\ even, \\ 5^{-5}(5X+1-4\sqrt{r})(5X+1+\sqrt{r}), & if sm/2\ odd. \end{array}$$

(ii) If $p \equiv 2(\mathrm{mod}\ 5)$ or $p \equiv 3(\mathrm{mod}\ 5)$, then

$$\psi_{(5,r)}(X) = \{ \begin{array}{ll} 5^{-5}(5X+1+4\sqrt{r})(5X+1-\sqrt{r}), & if sm/4\ even, \\ 5^{-5}(5X+1-4\sqrt{r})(5X+1+\sqrt{r}), & if sm/4\ odd. \end{array}$$

(iii) If $p \equiv 1(\mathrm{mod}\ 5)$, and $5 \nmid ms$, then $\psi_{(5,r)}(X)$ is irreducible over the rationals.

Lemma 2.5. *Let $N = 6$. We have the following results on the factorization of $\psi_{(6,r)}(X)$.*

(i) If $p \equiv 5(\mathrm{mod}\ 6)$, and ms is even, then

$$\psi_{(6,r)}(X) = \{ \begin{array}{ll} 6^{-6}(6X+1+5\sqrt{r})(6X+1-\sqrt{r}), & if sm/2\ even, \\ 6^{-6}(6X+1-5\sqrt{r})(6X+1+\sqrt{r}), & if sm/2\ odd. \end{array}$$

(ii) If $p \equiv 1(\mathrm{mod}\ 6)$,

(a) If $3 \nmid ms$, and ms is odd, then $\psi_{(6,r)}(X)$.is irreducible over the rationals.

(b) If $ms \equiv 0(\mathrm{mod}\ 6)$, then

$$\psi_{(6,r)}(X) = 6^{-6}[(6X+1) + (-1)^{\frac{kms}{2}} r^{\frac{1}{6}}(u+\overline{u}) + r^{\frac{1}{3}}(u_1+\overline{u}_1) + (-1)^{1+\frac{kms}{2}}\sqrt{r}]\times$$

$$[(6X+1) + (-1)^{\frac{kms}{2}} r^{\frac{1}{6}}(v+\overline{v}) + r^{\frac{1}{3}}(v_1+\overline{v}_1) + (-1)^{2+\frac{kms}{2}}\sqrt{r}]\times$$

$$\{(6X+1) + (-1)^{\frac{kms}{2}} r^{\frac{1}{6}}[(v+\overline{v}) - (u+\overline{u})] - r^{\frac{1}{3}}[(v_1+\overline{v}_1) + (u_1+\overline{u}_1]$$

$$+(-1)^{3+\frac{kms}{2}}\sqrt{r}\} \times [(6X+1) - (-1)^{\frac{kms}{2}} r^{\frac{1}{6}}(u+\overline{u}) + r^{\frac{1}{3}}(u_1+\overline{u}_1)+$$

$$(-1)^{4+\frac{kms}{2}}\sqrt{r}]\times[(6X+1)-(-1)^{\frac{kms}{2}} r^{\frac{1}{6}}(v+\overline{v})+r^{\frac{1}{3}}(v_1+\overline{v}_1)+(-1)^{5+\frac{kms}{2}}\sqrt{r}]$$

$$\times\{(6X+1) - (-1)^{\frac{kms}{2}} r^{\frac{1}{6}}[(v+\overline{v}) - (u+\overline{u})] - r^{\frac{1}{3}}[(v_1+\overline{v}_1) + (u_1+$$

$$\overline{u}_1] + (-1)^{6+\frac{kms}{2}}\sqrt{r}\}.$$

(c) If $ms \equiv 3(\mathrm{mod}\ 6)$, then

$$\psi_{(6,r)}(X) = 6^{-6}[(6X+1)^2 - 2r^{\frac{1}{3}}(u_1+\overline{u}_1)(6X+1) + p^{\frac{2}{3}ms}(u_1+\overline{u}_1)^2 - (-1)^k r^{\frac{1}{3}}(u+$$

$$\overline{u} - r^{\frac{1}{3}})^2] \times [(6X+1)^2 - 2r^{\frac{1}{3}}(v_1+\overline{v}_1)(6X+1) + p^{\frac{2}{3}ms}(v_1+\overline{v}_1)^2 -$$

$$(-1)^k r^{\frac{1}{3}}(v+\overline{v} - r^{\frac{1}{3}})^2] \times \{(6X+1)^2 + 2r^{\frac{1}{3}}[(v_1+\overline{v}_1) + (u_1+\overline{u}_1)](6X$$

$$+1) + p^{\frac{2}{3}ms}[(v_1+\overline{v}_1) + (u_1+\overline{u}_1)]^2 - (-1)^k r^{\frac{1}{3}}[(v+\overline{v}) - (u+\overline{u}) - r^{\frac{1}{3}})^2],$$

where u and v are given by $(u-v)(\overline{u}-\overline{v}) = p^{\frac{2}{3}ms}$, u_1 and v_1 are given by $(u+v)(\overline{u}+\overline{v}) = p^{\frac{1}{3}ms}$, and $p = 6k+1$.

Lemma 2.6. *Let $N = 7$.We have the following results on the factorization of $\psi_{(7,r)}(X)$.*

(i) If $p \equiv 6 (\mathrm{mod}\ 7)$, then

$$\psi_{(7,r)}(X) = \{ \begin{array}{ll} 7^{-7}(7X+1+6\sqrt{r})(7X+1-\sqrt{r}), & if sm/2\ even, \\ 7^{-7}(7X+1-6\sqrt{r})(7X+1+\sqrt{r}), & if sm/2\ odd. \end{array}$$

(ii) If $p \equiv 3 (\mathrm{mod}\ 7)$ or $p \equiv 5 (\mathrm{mod}\ 7)$, then

$$\psi_{(7,r)}(X) = \{ \begin{array}{ll} 7^{-7}(7X+1+6\sqrt{r})(7X+1-\sqrt{r}), & if sm/6\ even, \\ 7^{-7}(7X+1-6\sqrt{r})(7X+1+\sqrt{r}), & if sm/6\ odd. \end{array}$$

(iii) If $p \equiv 1 (\mathrm{mod}\ 7)$, and $7 \nmid ms$, then $\psi_{(7,r)}(X)$ is irreducible over the rationals.

Lemma 2.7. *Let $N = 8$. We have the following results on the factorization of $\psi_{(8,r)}(X)$.*

(i) If $p \equiv 7 (\mathrm{mod}\ 8)$, and ms is even, then

$$\psi_{(8,r)}(X) = \{ \begin{array}{ll} 8^{-8}(8X+1+7\sqrt{r})(8X+1-\sqrt{r}), & if sm/2\ even, \\ 8^{-8}(8X+1-7\sqrt{r})(8X+1+\sqrt{r}), & if sm/2\ odd. \end{array}$$

(ii) If $p \equiv 5 (\mathrm{mod}\ 8)$,

(a) If $ms \equiv 0 (\mathrm{mod}\ 8)$, then

$$\psi_{(8,r)}(X) = 8^{-8}[(8X+1) - \sqrt{r} + 8r^{\frac{1}{4}}cd]^2 \times [(8X+1) + 3\sqrt{r} - 4r^{\frac{1}{4}}c^2 + 8r^{\frac{3}{8}}d] \times$$

$$[(8X+1) - \sqrt{r} - 8r^{\frac{1}{4}}cd]^2 \times [(8X+1) - \sqrt{r} + 4r^{\frac{1}{4}}c^2 - 4r^{\frac{3}{8}}c] \times [(8X+$$

$$1) + 3\sqrt{r} - 4r^{\frac{1}{4}}c^2 - 8r^{\frac{3}{8}}d] \times [(8X+1) - \sqrt{r} + 4r^{\frac{1}{4}}c^2 + 4r^{\frac{3}{8}}c],$$

where c and d are given by $c^2 + 4d^2 = r^{\frac{1}{4}}, c \equiv 1 (\mathrm{mod}\ 4), \text{and } \mathrm{gcd(c, p)} = 1.$

(b) If $ms \equiv 4 (\mathrm{mod}\ 8)$, then

$$\psi_{(8,r)}(X) = 8^{-8}[(8X+1) - \sqrt{r} + 8r^{\frac{1}{4}}cd]^2 \times \{[(8X+1) + 3\sqrt{r} - 4r^{\frac{1}{4}}c^2]^2 - 64r^{\frac{3}{4}}d^2\}$$

$$\times [(8X+1) - \sqrt{r} - 8r^{\frac{1}{4}}cd]^2 \times \{[(8X+1) + 4r^{\frac{1}{4}}c^2 - \sqrt{r}]^2 - 16r^{\frac{1}{4}}c^2\},$$

where c and d are given by $c^2 + 4d^2 = r^{\frac{1}{4}}, c \equiv \pm 1 (\mathrm{mod}\ 4), \text{and } \mathrm{gcd(c, p)} = 1.$

(c) If $ms \equiv 2 (\mathrm{mod}\ 4)$, then

$$\psi_{(8,r)}(X) = 8^{-8}[(8X+1)^2 - 2\sqrt{r}(8X+1) + r - 16\sqrt{r}z^2]^2 \times [(8X+1)^4 + 4\sqrt{r}(8X$$

$$+1)^3 + 2\sqrt{r}(8X+1)^2(11\sqrt{r} - 4w^2) + 4r(8X+1)(9\sqrt{r} - 20w^2) +$$

$$r(9\sqrt{r} - 4w^2)^2],$$

where w and z are given by $w^2 + 4z^2 = \sqrt{r}, w \equiv 1 (\mathrm{mod}\ 4), \text{and } \mathrm{gcd(w, p)} = 1.$

(iii) If $p \equiv 3 (\mathrm{mod}\ 8)$,

(a) If $ms \equiv 0 (\mathrm{mod}\ 4)$, then

$$\psi_{(8,r)}(X) = 8^{-8}[(8X+1) - \sqrt{r}]^2 \times [(8X+1)^2 - 2\sqrt{r}(8X+1) + r - 16\sqrt{r}t^2]^2 \times$$

$$[(8X+1)^2 + 6\sqrt{r}(8X+1) + 9r - 16\sqrt{r}l^2],$$

where l *and* t *are given by* $l^2 + 2t^2 = \sqrt{r}, l \equiv -1 (\text{mod } 8), \text{and gcd(l, p)} = 1.$

(b) If $ms \equiv 2 (\text{mod } 4)$*, then*

$$\psi_{(8,r)}(X) = 8^{-8}[(8X+1)^2 + 2\sqrt{r}(8X+1) + r + 16\sqrt{r}t^2]^2 \times [(8X+1) - 3\sqrt{r}]^2$$
$$\times [(8X+1)^2 + 2\sqrt{r}(8X+1) + r + 16\sqrt{r}l^2],$$

where l *and* t *are given by* $l^2 + 2t^2 = \sqrt{r}, l \equiv \pm 1 (\text{mod } 4), \text{and gcd(l, p)} = 1.$

(iv) If $p \equiv 1 (\text{mod } 8)$,

(a) If ms *is odd, then* $\psi_{(8,r)}(X)$*.is irreducible over the rationals.*

(b) If $ms \equiv 0 (\text{mod } 8)$*, then*

$$\psi_{(8,r)}(X) = 8^{-8}[(8X+1) + \sqrt{r} + 8r^{\frac{1}{4}}cd + 2r^{\frac{1}{8}}(2c-4d)t] \times [(8X+1) + \sqrt{r} -$$
$$4r^{\frac{1}{4}}c^2 + 8r^{\frac{1}{8}}dl] \times [(8X+1) + \sqrt{r} - 8r^{\frac{1}{4}}cd + 2r^{\frac{1}{8}}(2c+4d)t] \times [(8X+$$
$$1) - 3\sqrt{r} + 4r^{\frac{1}{4}}c^2 - 4r^{\frac{1}{8}}cl] \times [(8X+1) + \sqrt{r} + 8r^{\frac{1}{4}}cd + 2r^{\frac{1}{8}}(4d-2c)t]$$
$$[(8X+1) + \sqrt{r} - 4r^{\frac{1}{4}}c^2 - 8r^{\frac{1}{8}}dl] \times [(8X+1) + \sqrt{r} - 8r^{\frac{1}{4}}cd - 2r^{\frac{1}{8}}(2c+4d)t]$$
$$\times [(8X+1) - 3\sqrt{r} + 4r^{\frac{1}{4}}c^2 + 4r^{\frac{1}{8}}cl],$$

where c *and* d *are given by* $c^2 + 4d^2 = r^{\frac{1}{4}}, c \equiv 1 (\text{mod } 8), \text{and gcd(c, p)} = 1;$ l *and* t *are given by* $l^2 + 2t^2 = \sqrt{r}, l \equiv 1 (\text{mod } 8), \text{and gcd(l, p)} = 1.$

(c) If $ms \equiv 4 (\text{mod } 8)$*, then*

$$\psi_{(8,r)}(X) = 8^{-8}[(8X+1)^2 + 2r^{\frac{1}{4}}(r^{\frac{1}{4}} + 8cd)(8X+1) + \sqrt{r}(r^{\frac{1}{4}} + 8cd)^2 - 4r^{\frac{1}{4}}(2c-$$
$$4d)^2t^2] \times [(8X+1)^2 + 2r^{\frac{1}{4}}(3r^{\frac{1}{4}} - 4c^2)(8X+1) + \sqrt{r}(3r^{\frac{1}{4}} - 4c^2)^2 - 64r^{\frac{1}{4}}d^2l^2]$$
$$\times [(8X+1)^2 + 2r^{\frac{1}{4}}(r^{\frac{1}{4}} - 8cd)(8X+1) + \sqrt{r}(r^{\frac{1}{4}} - 8cd)^2 - 4r^{\frac{1}{4}}(2c+4d)^2t^2] \times$$
$$[(8X+1)^2 + 2r^{\frac{1}{4}}(4c^2 - r^{\frac{1}{4}})(8X+1) + \sqrt{r}(4c^2 - r^{\frac{1}{4}})^2 - 16r^{\frac{1}{4}}c^2l^2,$$

where c *and* d *are given by* $c^2 + 4d^2 = r^{\frac{1}{4}}, c \equiv 1 (\text{mod } 4), \text{and gcd(c, p)} = 1;$ l *and* t *are given by* $l^2 + 2t^2 = \sqrt{r}, l \equiv 1 (\text{mod } 8), \text{and gcd(l, p)} = 1.$

(d) If $ms \equiv 2 (\text{mod } 4)$*, then*

$$\psi_{(8,r)}(X) = 8^{-8}\{(8X+1)^4 - 4\sqrt{r}(8X+1)^3 - 2\sqrt{r}(8X+1)^2(16z^2 + 16t^2 - 3\sqrt{r}) -$$
$$4\sqrt{r}(8X+1)[r - \sqrt{r}(16z^2 + 16t^2) + 128t^2z^2] + r(\sqrt{r} + 16z^2 - 16t^2)^2 -$$
$$64\sqrt{r}(\sqrt{r} - 16t^2)^2z^2\} \times (8X+1)^4 + 4\sqrt{r}(8X+1)^3 - 2\sqrt{r}(8X+1)^2$$
$$(4w^2 + 8l^2 - 3\sqrt{r}) + 4\sqrt{r}(8X+1)[r - \sqrt{r}(4w^2 + 8l^2) + 16w^2l^2] + r(\sqrt{r}$$
$$+4w^2 - 8l^2)^2 - 4\sqrt{r}(2\sqrt{r} - 4l^2)^2w^2,$$

where w *and* z *are given by* $w^2 + 4z^2 = \sqrt{r}, w \equiv 1 (\text{mod } 4), \text{and gcd(w, p)} = 1;$ l *and* t *are given by* $l^2 + 2t^2 = \sqrt{r}, l \equiv -1 (\text{mod } 4), \text{and gcd(l, p)} = 1.$

3. The weight enumerator of some classes of irreducible cyclic codes

Recall that $q = p^s$and $r = q^m$,where s and m are positive integers. $r - 1 = nN$ throughout this paper. Let $Z(r, a)$ denote the number of solutions $x \in F_r$ of the equation $Tr_{r/q}(ax^N) = 0$. We have then

$$Z(r,a) = \frac{1}{q}\sum_{x\in F_r}\sum_{\chi\in F_q^{\wedge}} \chi(Tr_{r/q}(ax^N)\overline{\chi(0)})$$

$$= \frac{1}{q}\sum_{x\in F_r}\sum_{y\in F_q} \chi_y(Tr_{r/q}(ax^N))$$

$$= \frac{1}{q}\sum_{x\in F_r}\sum_{y\in F_q} \chi_1(yTr_{r/q}(ax^N))$$

$$= \frac{1}{q}\sum_{x\in F_r}\sum_{y\in F_q} \chi_1(Tr_{r/q}(yax^N))$$

$$= \frac{1}{q}\sum_{x\in F_r}\sum_{y\in F_q} \chi_{r/p}(yax^N)$$

$$= \frac{1}{q}[r + q - 1 + \sum_{x\in F_{r^*}}\sum_{y\in F_{q^*}} \chi_1(Tr_{r/q}(yax^N))]$$

$$= \frac{1}{q}[r + q - 1 + N\sum_{y\in F_{q^*}}\sum_{x\in C_0^{(N,r)}} \chi_1(Tr_{r/q}(yax^N))]. \qquad (4)$$

Hence, for any $\beta \in F_r^*$ the Hamming weight of the codeword

$$\mathbf{c}(\beta) = (\mathrm{Tr}_{r/q}(\beta), \mathrm{Tr}_{r/q}(\beta\theta), \cdots, \mathrm{Tr}_{r/q}(\beta\theta^{n-1}))$$

in the ICC of (2) is equal to $n - \frac{Z(r,\beta)-1}{N}$.

Theorem 3.1. *([9]) If* $q \equiv 1(\mathrm{mod\ N})$ *and* $\gcd(\mathrm{m}, \mathrm{N}) = 1$, *then the set* $C(r, N)$ *in (2) is a* $[(q^m - 1)/N, m, q^{m-1}(q-1)/N]$ *code with the only nonzero weight* $q^{m-1}(q-1)/N$.

A. The weight enumerator in the Case $N = 5$

Lemma 3.2. *Let* $N = 5$. *If* $q \equiv 1(\mathrm{mod}\ 5)$, *then*

$$(r-1)/(q-1)\mathrm{mod\ N} = \mathrm{m\ mod\ 5}.$$

Proof. *Note that* $(r - 1)/(q - 1)\mathrm{mod}\ 5 = (\mathrm{q}^{\mathrm{m}-1} + \mathrm{q}^{\mathrm{m}-2} + \ldots + \mathrm{q} + 1)\mathrm{mod}\ 5$, *and* $q \equiv 1(\mathrm{mod}\ 5)$, *The conclusion then follows.* □

Theorem 3.3. *Let* $N = 5$ *and* $q \equiv 1 (\mathrm{mod}\ 5)$*. If* $5 \nmid m$*, then the set* $C(r, 5)$ *is a* $[(q^m - 1)/5, m, q^{m-1}(q - 1)/5]$ *code with the only nonzero weight* $q^{m-1}(q - 1)/5$*.*

Proof. Since gcd(m, 5) = 1. The conclusion then follows from Theorem 3.1. □

Example 3.4. *Let* $q = 11$ *and let* $m = 2$*. Then the set* $C(r, 5)$ *in (2) is a* $[24, 2, 22]$ *code with the only nonzero weight* 22*.*

Theorem 3.5. *Let* $q \equiv 1(\mathrm{mod}\ 5)$, $p \equiv 4(\mathrm{mod}\ 5)$*. and* $m \equiv 0(\mathrm{mod}\ 5)$*. Let* $r - 1 = nN$*, where*$N = 5$*.*

(i) If $sm \equiv 0(\mathrm{mod}\ 4)$*, then* $C(r, 5)$ *is an* $[(r - 1)/5, m, (q - 1)(r - \sqrt{r})/5q]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{4(r-1)}{5} x^{\frac{(q-1)(r-\sqrt{r})}{5q}} + \frac{(r-1)}{5} x^{\frac{(q-1)(r+4\sqrt{r})}{5q}}.$$

(ii) If $sm \equiv 2(\mathrm{mod}\ 4)$*, then* $C(r, 5)$ *is an* $[(r - 1)/5, m, (q - 1)(r - 4\sqrt{r})/5q]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{4(r-1)}{5} x^{\frac{(q-1)(r+\sqrt{r})}{5q}} + \frac{(r-1)}{5} x^{\frac{(q-1)(r-4\sqrt{r})}{5q}}.$$

Proof. If $q \equiv 1(\mathrm{mod}\ 5)$, $m \equiv 0(\mathrm{mod}\ 5)$, then by Lemma 3.2,$(r-1)/(q-1) \mathrm{mod}\ 5 = 0$. Let α be a fixed primitive element of F_r. Note that every element of F_q^* is the form $\alpha^{i(r-1)/(q-1)}$ for some integer i. Hence, $F_q^* \subset C_0^{(5,r)}$. It follows from Lemma 2.4 and (3) that the Gaussian period $\eta_0^{(5,r)} = \frac{-1-4\sqrt{r}}{5}$, $\eta_j^{(5,r)} = \frac{-1+\sqrt{r}}{5}$ $(1 \le j \le 4)$. Note that (4) becomes

$$Z(r, a) = \frac{1}{q}[r + q - 1 + N \sum_{y \in F_q^*} \sum_{x \in \in C_0^{(N,r)}} \chi(yax)]$$

$$= \frac{1}{q}[r + q - 1 + 5(q - 1)\eta_i^{(5,r)}],$$

where $a \in C_i^{(5,r)}$ for some i. Hence, the Hamming weight of the codeword $\mathbf{c}(\alpha)$ is

$$n - \frac{Z(r, a) - 1}{N} = \frac{r - Z(r, a)}{N} = \frac{(q - 1)(r - 1 - 5\eta_i^{(5,r)}}{5q}).$$

Then the weight enumerator can be determined in terms of (1).

The proof for the case that $sm \equiv 2(\mathrm{mod}\ 4)$ is the same as that of part(i). □

Theorem 3.6. *Let* $q \equiv 1(\mathrm{mod}\ 5)$, $p \equiv 2(\mathrm{mod}\ 5)$ *or* $p \equiv 3(\mathrm{mod}\ 5)$*. and* $m \equiv 0(\mathrm{mod}\ 5)$*. Let* $r - 1 = nN$*, where* $N = 5$*.*

(i) If $sm \equiv 0(\mathrm{mod}\ 8)$*, then* $C(r, 5)$ *is an* $[(r - 1)/5, m, (q - 1)(r - \sqrt{r})/5q]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{4(r-1)}{5} x^{\frac{(q-1)(r-\sqrt{r})}{5q}} + \frac{(r-1)}{5} x^{\frac{(q-1)(r+4\sqrt{r})}{5q}}.$$

(ii) If $sm \equiv 4 \pmod 8$, then $C(r,5)$ is an $[(r-1)/5, m, (q-1)(r-4\sqrt{r})/5q]$ code over F_q with the weight enumerator

$$1+\frac{4(r-1)}{5}x^{\frac{(q-1)(r+\sqrt{r})}{5q}}+\frac{(r-1)}{5}x^{\frac{(q-1)(r-4\sqrt{r})}{5q}}.$$

The proof of Theorem 3.6 is similar to that of the Theorem 3.5.
B. The weight enumerator in the Case $N = 6$

Lemma 3.7. *Let $N = 6$. If $q \equiv 1 \pmod 6$, then*

$$(r-1)/(q-1) \mathrm{mod\ N} = \mathrm{m\ mod\ 6}.$$

Theorem 3.8. *Let $N = 6$ and $q \equiv 1 \pmod 6$. If m is odd and $3 \nmid m$, then the set $C(r,6)$ is a $[(q^m-1)/6, m, q^{m-1}(q-1)/6]$ code with the only nonzero weight $q^{m-1}(q-1)/6$.*

Example 3.9. *Let $q = 7$ and $m = 5$. Then the set $C(r,6)$ in (2) is a* $[2801, 5, 2401]$ *code with the only nonzero weight* 2401.

Theorem 3.10. *Let $q \equiv 1 \pmod 6$, $p \equiv 5 \pmod 6$, and let s be even.*

(i) If $m \equiv 0 \pmod 6$, then $C(r,6)$ is an $[(r-1)/6, m, (q-1)(r-\sqrt{r})/6q]$ code over F_q with the weight enumerator

$$1+\frac{5(r-1)}{6}x^{\frac{(q-1)(r-\sqrt{r})}{6q}}+\frac{(r-1)}{6}x^{\frac{(q-1)(r+5\sqrt{r})}{6q}}.$$

(ii) If $m \equiv 2 \pmod 6$, then $C(r,6)$ is an $[(r-1)/6, m, (q-1)(r-\sqrt{r})/6q]$ code over F_q with the weight enumerator

$$1+\frac{r-1}{2}x^{\frac{(q-1)(r-\sqrt{r})}{6q}}+\frac{r-1}{2}x^{\frac{(q-1)(r+\sqrt{r})}{6q}}.$$

Proof. (i) Since $q \equiv 1 \pmod 6$, $m \equiv 0 \pmod 6$, then by Lemma 3.6,$(r-1)/(q-1) \mathrm{mod}\ 6 = 0$. Then similar to the proof of Theorem 3.5.

(ii) Since $q \equiv 1 \pmod 6$, $m \equiv 2 \pmod 6$, then by Lemma 3.7,$(r-1)/(q-1) \mathrm{mod}\ 6 = 2$. Hence, $\mathrm{gcd(6, r-1)/(q-1) modmod\ 6)} = 2$. From[12], If r be an even power of an odd prime, then

$$\chi(C_0^{(2,r)}) = \frac{-1\pm\sqrt{r}}{2}, \quad \chi(C_1^{(2,r)}) = \frac{-1\mp\sqrt{r}}{2}.$$

It then follows from (4) that

$$Z(r,a) = \frac{1}{q}[r+q-1+6\sum_{y\in F_q^*}\sum_{x\in\in C_0^{(6,r)}}\chi(yax)]$$

$$= \frac{1}{q}[r+q-1+6\times\frac{q-1}{3}\sum_{x\in C_0^{(2,r)}}\chi(ax)]$$

$$= \frac{1}{q}[r + q - 1 + (q-1)(-1 \pm \sqrt{r})].$$

Hence, the Hamming weight of the codeword $c(a)$ is

$$n - \frac{Z(r,a) - 1}{N} = \frac{r - Z(r,A)}{N} = \frac{(q-1)(r \pm \sqrt{r})}{6q}.$$

This is a complete proof for the second part. $\square$

Example 3.11. *Let* $q = 5^2$ *and* $m = 2$. *Then the set* $(r, 5)$ *in (2) is a* $[2801, 5, 2401]$ *code over* F_{25} *with the weight distribution* $1 + 312x^{96} + 312x^{104}$.

Theorem 3.12. *Let* $q \equiv 1(\text{mod } 6)$, $p \equiv 1(\text{mod } 6)$. .

(i) If $m \equiv 0(\text{mod } 6)$, *then* $C(r, 6)$ *is an* $[(r-1)/6, m]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{r-1}{6} x^{\frac{(q-1)[r+(-1)^{\frac{kms}{2}} r^{\frac{1}{6}}(u+\overline{u})+r^{\frac{1}{3}}(u_1+\overline{u}_1)+(-1)^{1+\frac{kms}{2}}\sqrt{r}]}{6q}} +$$

$$\frac{r-1}{6} x^{\frac{(q-1)[r+(-1)^{\frac{kms}{2}} r^{\frac{1}{6}}(v+\overline{v})+r^{\frac{1}{3}}(v_1+\overline{v}_1)+(-1)^{2+\frac{kms}{2}}\sqrt{r}]}{6q}} +$$

$$\frac{r-1}{6} x^{\frac{(q-1)\{r+(-1)^{\frac{kms}{2}} r^{\frac{1}{6}}[(v+\overline{v})-(u+\overline{u})]-r^{\frac{1}{3}}[(v_1+\overline{v}_1)+(u_1+\overline{u}_1)]+(-1)^{3+\frac{kms}{2}}\sqrt{r}\}}{6q}} +$$

$$\frac{r-1}{6} x^{\frac{(q-1)[r-(-1)^{\frac{kms}{2}} r^{\frac{1}{6}}(u+\overline{u})+r^{\frac{1}{3}}(u_1+\overline{u}_1)+(-1)^{4+\frac{kms}{2}}\sqrt{r}}{6q}]} +$$

$$\frac{r-1}{6} x^{\frac{(q-1)[r-(-1)^{\frac{kms}{2}} r^{\frac{1}{6}}(v+\overline{v})+r^{\frac{1}{3}}(v_1+\overline{v}_1)+(-1)^{5+\frac{kms}{2}}\sqrt{r}]}{6q}} +$$

$$\frac{r-1}{6} x^{\frac{(q-1)\{r-(-1)^{\frac{kms}{2}} r^{\frac{1}{6}}[(v+\overline{v})-(u+\overline{u})]-r^{\frac{1}{3}}[(v_1+\overline{v}_1)+(u_1+\overline{u}_1]+(-1)^{6+\frac{kms}{2}}\sqrt{r}\}}{6q}},$$

where u *and* v *are given by* $(u - v)(\overline{u} - \overline{v}) = p^{\frac{2}{3}ms}$, u_1 *and* v_1 *are given by* $(u + v)(\overline{u} + \overline{v})) = p^{\frac{1}{3}ms}$, *and* $p = 6k + 1$.

(ii) If $m \equiv 2(\text{mod } 6)$, *then* $C(r, 6)$ *is an* $[(r-1)/6, m, (q-1)(r - \sqrt{r})/6q]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{r-1}{2} x^{\frac{(q-1)(r-\sqrt{r})}{6q}} + \frac{r-1}{2} x^{\frac{(q-1)(r+\sqrt{r})}{6q}}.$$

The proof of Theorem 3.12 is similar to that of the Theorem 3.10.

Example 3.13. *Let $q = 7$ and $m = 2$. Then the set $C(r, 6)$ in (2) is a $[8, 2, 6]$ code over F_7 with the weight distribution $1 + 24x^6 + 24x^8$. We can then obtain nonzero codewords of $C(r, 6)$ in this example through programming, as follows :*

$\{4,4,5,2,3,3,2,5\}$, $\{6,3,6,0,1,4,1,0\}$, $\{4,6,6,4,3,1,1,3\}$, $\{0,2,1,2,0,5,6,5\}$,
$\{1,6,2,2,6,1,5,5\}$, $\{4,0,3,5,3,0,4,2\}$, $\{4,5,2,3,3,2,5,4\}$, $\{3,6,0,1,4,1,0,6\}$,
$\{6,6,4,3,1,1,3,4\}$, $\{2,1,2,0,5,6,5,0\}$, $\{6,2,2,6,1,5,5,1\}$, $\{0,3,5,3,0,4,2,4\}$,
$\{5,2,3,3,2,5,4,4\}$, $\{6,0,1,4,1,0,6,3\}$, $\{6,4,3,1,1,3,4,6\}$, $\{1,2,0,5,6,5,0,2\}$,
$\{2,2,6,1,5,5,1,6\}$, $\{3,5,3,0,4,2,4,0\}$, $\{2,3,3,2,5,4,4,5\}$, $\{0,1,4,1,0,6,3,6\}$,
$\{4,3,1,1,3,4,6,6\}$, $\{2,0,5,6,5,0,2,1\}$, $\{2,6,1,5,5,1,6,2\}$, $\{5,3,0,4,2,4,0,3\}$,
$\{3,3,2,5,4,4,5,2\}$, $\{1,4,1,0,6,3,6,0\}$, $\{3,1,1,3,4,6,6,4\}$, $\{0,5,6,5,0,2,1,2\}$,
$\{6,1,5,5,1,6,2,2\}$, $\{3,0,4,2,4,0,3,5\}$, $\{3,2,5,4,4,5,2,3\}$, $\{4,1,0,6,3,6,0,1\}$,
$\{1,1,3,4,6,6,4,3\}$, $\{5,6,5,0,2,1,2,0\}$, $\{1,5,5,1,6,2,2,6\}$, $\{0,4,2,4,0,3,5,3\}$,
$\{2,5,4,4,5,2,3,3\}$, $\{1,0,6,3,6,0,1,4\}$, $\{1,3,4,6,6,4,3,1\}$, $\{6,5,0,2,1,2,0,5\}$,
$\{5,5,1,6,2,2,6,1\}$, $\{4,2,4,0,3,5,3,0\}$, $\{5,4,4,5,2,3,3,2\}$, $\{0,6,3,6,0,1,4,1\}$,
$\{3,4,6,6,4,3,1,1\}$, $\{5,0,2,1,2,0,5,6\}$, $\{5,1,6,2,2,6,1,5\}$, $\{2,4,0,3,5,3,0,4\}$,

It is obvious that the number of nonzero codewords with Hamming weight either 6 or 8 is 24.

C. The weight enumerator in the Case $N = 7$

Lemma 3.14. *Let $N = 7$. If $q \equiv 1 (\mathrm{mod}\ 7)$, then*

$$(r-1)/(q-1) \mathrm{mod\ N} = \mathrm{m\ mod\ 7}.$$

Theorem 3.15. *Let $N = 7$, $q \equiv 1 (\mathrm{mod}\ 7)$, and $7 \nmid m$. Then the set $C(r, 7)$ is a $[(q^m - 1)/7, m, q^{m-1}(q - 1)/7]$ code with the only nonzero weight $q^{m-1}(q - 1)/7$.*

Example 3.16. *Let $q = 29$ and $m = 2$. Then the set $C(r, 7)$ in (2) is a $[120, 2, 116]$ code with the only nonzero weight 116.*

Theorem 3.17. *Let $q \equiv 1 (\mathrm{mod}\ 7)$, $p \equiv 6 (\mathrm{mod}\ 7)$, and $m \equiv 0 (\mathrm{mod}\ 7)$. Let $r - 1 = nN$, where $N = 7$.*

(i) If $sm \equiv 0 (\mathrm{mod}\ 4)$, then $C(r, 7)$ is an $[(r-1)/7, m, (q-1)(r-\sqrt{r})/7q]$ code over F_q with the weight enumerator

$$1 + \frac{6(r-1)}{7} x^{\frac{(q-1)(r-\sqrt{r})}{7q}} + \frac{(r-1)}{7} x^{\frac{(q-1)(r+6\sqrt{r})}{7q}}.$$

(ii) If $sm \equiv 2 (\mathrm{mod}\ 4)$, then $C(r, 7)$ is an $[(r-1)/7, m, (q-1)(r-6\sqrt{r})/7q]$ code over F_q with the weight enumerator

$$1 + \frac{6(r-1)}{7} x^{\frac{(q-1)(r+\sqrt{r})}{7q}} + \frac{(r-1)}{7} x^{\frac{(q-1)(r-6\sqrt{r})}{7q}}.$$

Theorem 3.18. *Let* $q \equiv 1 (\mathrm{mod}\ 7)$, $p \equiv 3 (\mathrm{mod}\ 7)$ *or* $p \equiv 5 (\mathrm{mod}\ 7)$ *and* $m \equiv 0 (\mathrm{mod}\ 7)$. *Let* $r - 1 = nN$, *where* $N = 7$.

(i) If $sm \equiv 0 (\mathrm{mod}\ 12)$, *then* $C(r, 7)$ *is an* $[(r-1)/7, m, (q-1)(r-\sqrt{r})/7q]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{6(r-1)}{7} x^{\frac{(q-1)(r-\sqrt{r})}{7q}} + \frac{(r-1)}{7} x^{\frac{(q-1)(r+6\sqrt{r})}{7q}}.$$

(ii) If $sm \equiv 6 (\mathrm{mod}\ 12)$, *then* $C(r, 7)$ *is an* $[(r-1)/7, m, (q-1)(r-6\sqrt{r})/7q]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{6(r-1)}{7} x^{\frac{(q-1)(r+\sqrt{r})}{7q}} + \frac{(r-1)}{7} x^{\frac{(q-1)(r-6\sqrt{r})}{7q}}.$$

The proof of Theorem 3.17, 3.18 is similar to that of the Theorem 3.5.

From[13], We have the following theorem.

Theorem 3.19. *Let* $N = 7$, $r = q^{3m}$, $p = q$ *and* $a = \frac{\omega_p(n)}{p-1}$. *Let* $n = n_0 + n_1 p + \cdots$ *be the expansion of* n *in the base* p *and let* $\omega_p(n) = n_0 + n_1 + \cdots$. *Let* c_m, d_m *be the unique positive integers prime to* p *that satisfy the diophantine equation* $c_m^2 + N d_m^2 = 4p^{m(3-2a)}$. *Then* $C(r, 7)$ *of length* n_m *have three weights with the weight distribution*

$$1 + n_m x^{\frac{(p-1)(2r \pm 3p^{ma} c_m)}{7p}} + 3n_m x^{\frac{(p-1)[2r \pm p^{ma}(7d_m - c_m)]}{14p}} + 3n_m x^{\frac{(p-1)[2r \mp p^{ma}(7d_m + c_m)]}{14p}}. \tag{5}$$

Example 3.20. *Let* $p = 2$ *and let* $m = 2$. *Then* $n_2 = 9, a = d_2 = 1, c_2 = 3$. *It then follows from (2) that* $C(r, 7)$ *in (2) is a* $[9, 2, 2]$ *code over* F_2 *with the weight distribution* $1 + 9x^2 + 27x^4 + 27x^6$. *We can then obtain all codewords of* $C(r, 7)$ *in this example through programming, as follows :*

$\{1,1,0,0,1,0,1,0,0\}, \{1,0,1,1,0,0,0,0,1\}, \{1,1,1,1,0,1,0,1,0\}, \{1,0,0,0,1,0,1,1,0\},$
$\{1,1,1,1,1,1,0,0,0\}, \{1,1,1,0,1,1,1,0,0\}, \{0,0,1,0,0,1,0,0,0\}, \{1,0,0,1,0,1,0,0,1\},$
$\{0,1,1,0,0,0,0,1,1\}, \{1,1,1,0,1,0,1,0,1\}, \{0,0,0,1,0,1,1,0,1\}, \{1,1,1,1,1,0,0,0,1\},$
$\{1,1,0,1,1,1,0,0,1\}, \{1,1,0,1,1,0,0,0,0\}, \{0,1,1,1,1,0,1,0,1\}, \{0,1,0,0,0,1,0,1,1\},$
$\{0,1,0,0,1,0,0,0,0\}, \{0,0,1,0,1,0,0,1,1\}, \{1,1,0,0,0,0,1,1,0\}, \{1,1,0,1,0,1,0,1,1\},$
$\{0,0,1,0,1,1,0,1,0\}, \{1,1,1,1,0,0,0,1,1\}, \{1,0,1,1,1,0,0,1,1\}, \{1,0,0,1,0,0,0,0,0\},$
$\{0,1,0,1,0,0,1,1,0\}, \{1,0,0,0,0,1,1,0,1\}, \{1,0,1,0,1,0,1,1,1\}, \{0,1,0,1,1,0,1,0,0\},$
$\{1,1,1,0,0,0,1,1,1\}, \{0,1,1,1,0,0,1,1,1\}, \{0,0,1,0,0,0,0,0,1\}, \{1,0,1,0,0,1,1,0,0\},$
$\{0,0,0,0,1,1,0,1,1\}, \{0,1,0,1,0,1,1,1,1\}, \{1,0,1,1,0,1,0,0,0\}, \{1,1,0,0,0,1,1,1,1\},$
$\{1,1,1,0,0,1,1,1,0\}, \{0,1,0,0,0,0,0,1,0\}, \{0,1,0,0,1,1,0,0,1\}, \{0,0,0,1,1,0,1,1,0\},$
$\{1,0,1,0,1,1,1,1,0\}, \{0,1,1,0,1,0,0,0,1\}, \{1,0,0,0,1,1,1,1,1\}, \{1,1,0,0,1,1,1,0,1\},$
$\{1,0,0,0,0,0,1,0,0\}, \{1,0,0,1,1,0,0,1,0\}, \{0,0,1,1,0,1,1,0,0\}, \{0,1,0,1,1,1,1,0,1\},$
$\{1,1,0,1,0,0,0,1,0\}, \{0,0,0,1,1,1,1,1,1\}, \{1,0,0,1,1,1,0,1,1\}, \{0,0,0,0,0,1,0,0,1\},$
$\{0,0,1,1,0,0,1,0,1\}, \{0,1,1,0,1,1,0,0,0\}, \{1,0,1,1,1,1,0,1,0\}, \{1,0,1,0,0,0,1,0,1\},$
$\{0,0,1,1,1,1,1,1,0\}, \{0,0,1,1,1,0,1,1,1\}, \{0,0,0,0,1,0,0,1,0\}, \{0,1,1,0,0,1,0,1,0\},$
$\{0,1,1,1,1,1,1,0,0\}, \{0,1,1,1,0,1,1,1,0\}, \{0,0,0,1,0,0,1,0,0\}, \{0,0,0,0,0,0,0,0,0\},$

It is obvious that the number of nonzero codewords with Hamming weight $2, 4$ *or* 6 *is* $9, 27$ *or* 27, *respectively.*

D. The weight enumerator in the Case $N = 8$

Lemma 3.21. *Let* $N = 8$. *If* $q \equiv 1(\text{mod}8)$, *then*

$$(r-1)/(q-1)\text{mod N} = \text{m mod 8}.$$

Theorem 3.22. *Let* $N = 8$ *and* $q \equiv 1(\text{mod } 8)$. *If* m *is odd , then the set* $C(r, 8)$ *is a* $[(q^m - 1)/8, m, q^{m-1}(q-1)/8]$ *code with the only nonzero weight* $q^{m-1}(q-1)/8$.

Example 3.23. *Let* $q = 17$ *and let* $m = 3$. *Then the set* $C(r, 8)$ *in (2) is a* $[614, 3, 578]$ *code with the only nonzero weight 578.*

Theorem 3.24. *Let* $q \equiv 1(\text{mod } 8)$, $p \equiv 7(\text{mod } 8)$, *and* s *be even.*

(i) If $m \equiv 0(\text{mod } 8)$, *then* $C(r, 8)$ *is an* $[(r-1)/8, m, (q-1)(r-\sqrt{r})/8q]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{7(r-1)}{8} x^{\frac{(q-1)(r-\sqrt{r})}{8q}} + \frac{(r-1)}{8} x^{\frac{(q-1)(r+7\sqrt{r})}{8q}}.$$

(ii) If $m \equiv 2(\text{mod } 8)$, *then* $C(r, 8)$ *is an* $[(r-1)/6, m, (q-1)(r-\sqrt{r})/8q]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{r-1}{2} x^{\frac{(q-1)(r-\sqrt{r})}{8q}} + \frac{r-1}{2} x^{\frac{(q-1)(r+\sqrt{r})}{8q}}.$$

The proof of Theorem 3.24 is similar to that of the Theorem 3.10.

Theorem 3.25. *Let* $q \equiv 1(\text{mod } 8)$, $p \equiv 5(\text{mod } 8)$, *and let* s *be even.*

(i) If $m \equiv 0(\text{mod } 8)$, *then* $C(r, 8)$ *is an* $[(r-1)/8, m]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{r-1}{4} x^{\frac{(q-1)(r-\sqrt{r}+8r^{\frac{1}{4}}cd)}{8q}} + \frac{r-1}{4} x^{\frac{(q-1)(r-\sqrt{r}-8r^{\frac{1}{4}}cd)}{8q}} + \frac{r-1}{8} x^{\frac{(q-1)(r+3\sqrt{r}-4r^{\frac{1}{4}}c^2+8r^{\frac{3}{8}}d)}{8q}} +$$

$$\frac{r-1}{8} x^{\frac{(q-1)(r-\sqrt{r}+4r^{\frac{1}{4}}c^2-4r^{\frac{3}{8}}c)}{8q}} + \frac{r-1}{8} x^{\frac{(q-1)(r+3\sqrt{r}-4r^{\frac{1}{4}}c^2-8r^{\frac{3}{8}}d)}{8q}} + \frac{r-1}{8} x^{\frac{(q-1)(r-\sqrt{r}+4r^{\frac{1}{4}}c^2+4r^{\frac{3}{8}}c)}{8q}},$$

where c *and* d *are given by* $c^2 + 4d^2 = r^{\frac{1}{4}}, c \equiv 1(\text{mod } 4), \text{and } \gcd(\text{c}, \text{p}) = 1$.

(ii) If $m \equiv 2(\text{mod } 8)$, *then* $C(r, 8)$ *is an* $[(r-1)/8, m, (q-1)(r-\sqrt{r})/8q]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{r-1}{2} x^{\frac{(q-1)(r-\sqrt{r})}{8q}} + \frac{r-1}{2} x^{\frac{(q-1)(r+\sqrt{r})}{8q}}.$$

The proof of Theorem 3.25 is similar to that of the Theorem 3.10.

Theorem 3.26. *Let* $q \equiv 1 (\text{mod } 8)$, $p \equiv 3 (\text{mod } 8)$. *and let* s *be even. If* $m \equiv 2 (\text{mod } 8)$, *then* $C(r, 8)$ *is an* $[(r-1)/8, m, (q-1)(r-\sqrt{r})/8q]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{r-1}{2} x^{\frac{(q-1)(r-\sqrt{r})}{8q}} + \frac{r-1}{2} x^{\frac{(q-1)(r+\sqrt{r})}{8q}}.$$

The proof of Theorem 3.26 is similar to that of the Theorem 3.10(ii).

Example 3.27. *Let* $q = 3^2$ *and let* $m = 2$. *Then* $C(r, 8)$ *in (2) is a* $[10, 2, 8]$ *code over* F_9 *with the weight distribution* $1 + 40x^8 + 40x^{10}$.

Theorem 3.28. *Let* $q \equiv 1 (\text{mod } 8)$, $p \equiv 1 (\text{mod } 8)$.

(i) If $m \equiv 0 (\text{mod } 8)$, *then* $C(r, 8)$ *is an* $[(r-1)/8, m]$ *code over* F_q *with the weight enumerator*

$$\begin{aligned} &1 + \frac{r-1}{8} x^{\frac{(q-1)[r+\sqrt{r}+8r^{\frac{1}{4}}cd+2r^{\frac{1}{8}}(2c-4d)t]}{8q}} + \frac{r-1}{8} x^{\frac{(q-1)[r+\sqrt{r}-4r^{\frac{1}{4}}c^2+8r^{\frac{1}{8}}dl]}{8q}} + \\ &\frac{r-1}{8} x^{\frac{(q-1)[r+\sqrt{r}-8r^{\frac{1}{4}}cd+2r^{\frac{1}{8}}(2c+4d)t]}{8q}} + \frac{r-1}{8} x^{\frac{(q-1)[r-3\sqrt{r}+4r^{\frac{1}{4}}c^2-4r^{\frac{1}{8}}cl]}{8q}} + \\ &\frac{r-1}{8} x^{\frac{(q-1)[r+\sqrt{r}+8r^{\frac{1}{4}}cd+2r^{\frac{1}{8}}(4d-2c)t]}{8q}} + \frac{r-1}{8} x^{\frac{(q-1)[r+\sqrt{r}-4r^{\frac{1}{4}}c^2-8r^{\frac{1}{8}}dl]}{8q}} + \\ &\frac{r-1}{8} x^{\frac{(q-1)[r+\sqrt{r}-8r^{\frac{1}{4}}cd-2r^{\frac{1}{8}}(2c+4d)t]}{8q}} + \frac{r-1}{8} x^{\frac{(q-1)[r-3\sqrt{r}+4r^{\frac{1}{4}}c^2+4r^{\frac{1}{8}}cl]}{8q}}, \end{aligned}$$

where c *and* d *are given by* $c^2 + 4d^2 = r^{\frac{1}{4}}, c \equiv 1 (\text{mod } 8), \text{andgcd}(\text{c}, \text{p}) = 1$; l *and* t *are given by* $l^2 + 2t^2 = \sqrt{r}, l \equiv 1 (\text{mod } 8), \text{and } \gcd(\text{l}, \text{p}) = 1$.

(ii) If $m \equiv 2 (\text{mod } 8)$, *then* $C(r, 8)$ *is an* $[(r-1)/8, m, (q-1)(r-\sqrt{r})/8q]$ *code over* F_q *with the weight enumerator*

$$1 + \frac{r-1}{2} x^{\frac{(q-1)(r-\sqrt{r})}{8q}} + \frac{r-1}{2} x^{\frac{(q-1)(r+\sqrt{r})}{8q}}.$$

The proof of Theorem 3.28 is similar to that of the Theorem 3.10.

Example 3.29. *Let* $q = 17$ *and let* $m = 2$. *Then* $C(r, 8)$ *in (2) is a* $[36, 2, 32]$ *code over* F_{17} *with the weight distribution* $1 + 144x^{32} + 144x^{36}$.

4. Conclusions

In this paper, we studied the weight enumerator of irreducible cyclic codes $C(r, N)$ for all $5 \leq N \leq 8$, and gave some specific examples. Furthermore, all codewords of $C(r, N)$ in certain examples can be obtained through programming, which is beneficial to verify the weight distributions of codes. Due to space limitation and the complexity of the weight enumerator of codes, it is hard to describe the weight distributions in more general cases. We shall work on this subject in a future work.

5. Acknowledgements

This work was supported by the National Natural Science Foundation of China (Grant No. 60873119).

6. References

References

[1] C. Ding, D. Feng , C. Ma, et al.. The weight enumerator of a class of cyclic codes[J]. IEEE Trans. Inform. Theory, 2011,57(1): 397-402.

[2] J.Yuan, C.Carlet, and C.Ding. The weight distribution of a class of linear codes from perfect nonlinear functions[J]. IEEE Trans. Inform. Theory, 2006,52(2): 712-717.

[3] Z.H. Li, T. Xue, and H. Lai. Secret sharing schemes from binary linear codes[J]. Information Science, 2011,180(22): 4412-4419.

[4] R.J.McEliece and H. Rumsey Jr.. Euler produces, cyclotomy, and coding[J]. Number Theroy,1972, 4:302-311.

[5] B.Schmidt and C.White. All two-weight irreducible cyclic codes? [J]. Finite Fields Appl., 2002, 8:1-17.

[6] P.Langevin. A new class of two weight codes[J].. Finite Fields Appl., 1996,181-187.

[7] L.D.Baumert and R.J.McEliece. Weights of irreducible cyclic codes[J]. Inf.Contr., 1972,20(2):158-175.

[8] L.D.Baumert and J.Mykkeltveit. Weight distributions of some irreducible cyclic codes[J]. DSN Progr. Rep., 1973,16:128-131.

[9] C. Ding . The weight distribution of some irreducible cyclic codes[J]. IEEE Trans. Inform. Theory, 2009, 55 (3): 955-960.

[10] S.Gurak. Period polynomials for of fixed small degree[J]. Number Theroy, 2004,127-145.

[11] A.Hoshi. Explicit lifts of quintic Jacobi sums and period polynomials for [J]. Proc. Japan Acad., 2006, 82(7):87-92.

[12] C. Ding . Complex codebooks from combinatorial designs. IEEE Trans. Inform. Theory, 2006,52(9): 4229-1235.

Zhi Hui Li
College of Mathematics and Information Science
Shaanxi Normal University
Xi'an 710062, P. R. China
e-mail:snnulzh@yahoo.com.cn